**Texila International Journal | Open Access | Peer Reviewed**

Journal: Texila International Journal of Academic Research
https://doi.org/10.21522/TIJAR.2014.13.03.Art019

ISSN: 2520-3088
Volume 13 Issue 3, 2026

# The Influence of Multi-stakeholder Health Promotion Interventions on Maternal and Infant Mortality Outcomes in Lagos State, Nigeria

**Bolajoko Esther Ogunwale**

*Department of Public Health, Texila American University, Georgetown, Guyana, South America*

**Correspondence to be addressed:** E-mail - bolajokoogunwale@gmail.com

**Cite this article as**: Ogunwale BE. The influence of multi-stakeholder health promotion interventions on maternal and infant mortality outcomes in Lagos State, Nigeria. Texila Int J Acad Res. 2026;13(3).
**https://doi.org/10.21522/TIJAR.2014.13.03.Art019**

**Received**: 21.05.2026
**Accepted**: 04.07.2026
**Published**: 31.07.2026

***Abstract*** *Maternal and infant mortality rates in peri-urban Lagos State, Nigeria, persist at critically elevated levels despite improvements in health infrastructure. In the Epe and Ibeju Lekki Local Government Areas (LGAs), a trust deficit rooted in sociocultural disconnection between formal health systems and community governance structures drives high rates of unbooked deliveries and late-stage obstetric emergencies. This study evaluates the influence of a structured Multi-Stakeholder Health Promotion Intervention (MSHPI) on maternal health service utilization, emphasizing the roles of traditional leaders, religious authorities, male household decision-makers, and narrative media in driving behavioural change. A convergent parallel mixed-methods design was employed. A structured questionnaire was administered to 500 women of reproductive age (15 to 49 years) across Epe and Ibeju Lekki LGAs using multi-stage stratified sampling, with data analyzed via logistic regression and Chi-square tests. Qualitative data from 25 Key Informant Interviews and 4 Focus Group Discussions were analyzed through thematic content analysis. Results revealed that women receiving antenatal care (ANC) counselling endorsed by traditional rulers were 3.42 times more likely to book care in the first trimester (OR = 3.42; 95% CI: 2.15, 5.43; $p < 0.001$). Husband's approval was the strongest individual predictor of skilled birth attendance (OR = 4.42; $p < 0.001$). Exposure to the 'Man of the House' edutainment intervention produced a 68.4% increase in facility-based deliveries ($p = 0.004$). Postnatal care uptake remained critically low at 29.6%. Integrating traditional, religious, and male stakeholders into a coordinated advocacy framework significantly improves health-seeking behaviour, highlighting the value of institutionalizing Community Health Advocacy Councils.*

***Keywords*** *Antenatal Care, Community Health Promotion, Edutainment, Infant Mortality, Lagos State, Maternal Mortality, Multi-stakeholder Interventions, Male Involvement, Nigeria, Trust Infrastructure.*

***This article is published under Creative Commons Attribution License (CC BY 4.0)***

https://doi.org/10.21522/TIJAR.2014.13.03.Art019

## Introduction

The global trajectory of public health is undergoing a decisive transformation, one that moves beyond the metrics of hospital bed density and surgical capacity toward a deeper reckoning with the social, cultural, and relational determinants of health outcomes. Nowhere is this transformation more urgently needed than in sub-Saharan Africa, where maternal and infant mortality rates remain among the highest in the world, and where the failure of conventional clinical strategies to reduce these rates has become undeniable [18, 19, 23].

Nigeria occupies a troubling position at the centre of this crisis. Despite being the continent's largest economy and most populous nation, Nigeria accounts for a disproportionate share of global maternal deaths. Within this national context, Lagos State presents a paradox that is both intellectually compelling and morally urgent. As the country's commercial capital, Lagos possesses the infrastructure, medical personnel, and financial resources that other states lack. Yet in its peripheral LGAs, particularly Epe and Ibeju Lekki, maternal and infant mortality rates persist at levels incompatible with this level of economic development [10, 18, 23].

This paradox demands explanation. The answer, this study argues, lies not in a shortage of facilities but in a chronic deficit of trust. Expectant mothers in these communities do not necessarily distrust medicine; they distrust a health system that has repeatedly failed to communicate with them on their own cultural terms, through the voices they consider authoritative. When a traditional ruler, an Imam, a Pastor, or a husband endorses a health behaviour, the message carries a social legitimacy that a clinical poster or a health talk at the Primary Health Centre (PHC) cannot replicate.

This study investigates the influence of Multi-Stakeholder Health Promotion Interventions (MSHPIs), which are structured frameworks that deliberately engage traditional leaders, religious authorities, male household decision-makers, and narrative media, on maternal and infant health outcomes in Epe and Ibeju Lekki. It does so through a mixed-methods design that combines the statistical rigour of quantitative analysis with the interpretive depth of qualitative inquiry, producing a body of evidence that is both numerically verifiable and contextually meaningful.

## Background

The 2024 Nigeria Verbal and Social Autopsy (VASA) survey identified that more than 60% of maternal deaths in the southwest geopolitical zone were precipitated by Type 1 Delays, which represent delays in the decision to seek care. These are not delays caused by road conditions or clinic opening hours; they are delays caused by the absence of social permission. A woman in Epe waits for her husband's approval. She waits for her mother-in-law's counsel. She waits for her pastor to confirm that hospital birth is spiritually permissible. When these approvals do not come on time, the consequences are fatal [11].

Retrospective analyses of Lagos tertiary facilities consistently demonstrate that unbooked mothers, meaning those who are present at labour without prior ANC contact, face mortality rates many times higher than their booked counterparts. The unbooked status itself is a symptom of the Trust Deficit: a system that has not earned the confidence of the communities it claims to serve [13, 15].

The Epe and Ibeju Lekki LGAs were selected as study sites because they represent the intersection of two realities: rapid economic growth and deeply embedded traditional community governance. These are not remote, resource-deprived localities. They are LGAs connected to Nigeria's most dynamic metropolitan economy, and yet they carry health burdens that mirror far more isolated communities. This makes them an ideal and

urgent case study for understanding what separates infrastructure from impact in Nigerian public health.

## Theoretical Framework

### The Social Ecological Model (SEM)

The Social Ecological Model provides structural architecture for this study. The SEM conceptualizes health behaviour as the outcome of interactions across five nested levels: individual, interpersonal, community, institutional, and policy. Applied to maternal health in peri-urban Lagos, the model reveals that interventions concentrated at the individual level, such as patient education and antenatal counselling, have limited reach when the interpersonal and community layers remain unconfigured for health promotion. In this study, the SEM is operationalized by treating community stakeholders, including traditional rulers, religious leaders, and male household heads, as the interpersonal and community layers through which clinical messages must pass in order to be acted upon. The model predicts, and this study confirms, that aligning these layers dramatically improves the penetration of health-promoting norms [9, 21].

### The Trust Infrastructure Theory

This study introduces the Trust Infrastructure Theory as an original theoretical contribution. Trust Infrastructure is defined as the network of relational validations, social endorsements, and cultural authority structures that determine whether a formal health system is perceived as credible, safe, and worth engaging with. In the absence of robust Trust Infrastructure, clinical facilities may be physically accessible yet socially inaccessible, meaning they are present in the community but absent from the community's confidence.

The theory posits that Trust Infrastructure is not built by the health system alone; it is borrowed from existing community authority systems. When a traditional ruler endorses the PHC, the PHC inherits a portion of the trust already deposited in that ruler over generations. This borrowed legitimacy is quantifiable, as the data in this study demonstrate.

### The Dual-Barricade Model

Factor analysis of the barriers identified in this study yields a novel framework: the Dual-Barricade Model, which classifies obstacles to maternal healthcare utilization into two distinct categories. Structural Barricades include transport difficulties, geographic distance, financial costs, and poor road infrastructure. Socio-Psychological Barricades include husband's disapproval, fear of medical procedures, cultural or religious beliefs, and misinformation regarding clinical interventions.

Critically, the data show that Socio-Psychological Barricades exert a stronger predictive influence on delivery outcomes than Structural Barricades. This finding fundamentally reorients the policy conversation: addressing roads and fees, while necessary, is insufficient. The inner barricade, representing psychological and relational resistance, must be dismantled first, and this requires the tools of community trust rather than clinical instruction.

## Methods

### Study Design

This study employs a Convergent Parallel Mixed-Methods Design. Quantitative and qualitative data were collected concurrently, analyzed independently, and integrated during interpretation. This design ensures methodological triangulation: statistical associations are validated and contextualized by the lived experiences and institutional knowledge of key community actors, producing a more complete and credible evidence base than either approach alone could achieve.

### Study Area

The study was conducted in Epe and Ibeju Lekki LGAs of Lagos State, selected for their

hybrid socio-demographic profile: economically transitioning peri-urban communities with retained traditional governance structures and documented disparities in maternal health outcomes. Both LGAs are served by Primary Health Centres under the Lagos State Primary Health Care Board (LSPHCB) [8].

### Study Population and Sampling

The quantitative sample comprised 500 women of reproductive age (15 to 49 years) who had experienced at least one pregnancy in the preceding 24 months. Sample size was determined using the Cochran formula at a 95% confidence level and 5% margin of error. A multi-stage stratified sampling technique was applied: purposive LGA selection, stratified ward selection, and systematic household sampling within wards.

The qualitative sample was purposively selected to include 25 key informants, comprising traditional rulers, senior religious leaders, PHC facility heads, and community health extension workers (CHEWs), and participants for four Focus Group Discussions (FGDs) composed of male household decision-makers and community women's groups.

### Instrumentation

Quantitative data were collected using the Multidimensional Maternal Health Questionnaire (MMHQ), a structured instrument validated through academic review and pilot-tested in a comparable peri-urban Lagos community. Internal consistency was confirmed at Cronbach's alpha = 0.78. The instrument collected data across five domains: socio-demographic profile, ANC utilization patterns, primary source of trusted health information, perceived barriers to PHC attendance, and male involvement in reproductive health decisions.

Qualitative data were gathered through semi-structured Key Informant Interview (KII) guides and FGD protocols, conducted in both English and Yoruba (Ijebu dialect). All sessions were audio-recorded with informed consent and transcribed verbatim.

### Data Analysis

Quantitative data was analyzed using IBM SPSS Version 28. Descriptive statistics characterize the sample profile. Binary logistic regression identified predictors of early ANC booking and skilled birth attendance, reported as Odds Ratios (OR) with 95% confidence intervals. Chi-square tests evaluated the significance of categorical associations, including the impact of edutainment exposure on male behaviour.

Qualitative transcripts were analyzed through thematic content analysis following Braun and Clarke's six-phase framework [3]. Key emergent themes included Trust Infrastructure, the Garrison Effect (male decision authority), Ancestral Compliance, and the Sovereignty of the Signal.

## Results

### Socio-demographic Profile

Of the 500 respondents, the majority (58.4%; n = 292) were aged 25 to 34 years, representing the peak reproductive window in this population. Most (72.8%; n = 364) were married and co-residing with their husbands, a finding that significantly contextualizes the role of male decision authority documented throughout the study. Secondary school completion was reported by 44.0%, with 31.2% holding post-secondary qualifications. Despite moderate educational attainment, 52.0% of households reported monthly incomes below ₦100,000, highlighting the economic vulnerability that constrains healthcare-seeking autonomy [10].

### The Maternal Care Continuum: From Entry to Dropout

Analysis of care-seeking patterns across the maternal health continuum reveals a progressive decline in service engagement from

https://doi.org/10.21522/TIJAR.2014.13.03.Art019

ANC initiation through to postnatal care, a pattern consistent with the hypothesis that barriers intensify as the pregnancy advances toward delivery. Comprehensive metrics documenting this attrition pattern across the different sequential stages of the pregnancy cycle are outlined in Table 1.

**Table 1.** Maternal Care Continuum and Service Leakage (N = 500)

| Stage of Maternal Care | n | % | Interpretation |
|---|---|---|---|
| ANC Initiation (≥1 visit) | 412 | 82.4% | High entry into care pathway |
| Optimal ANC Completion (4+ visits) | 314 | 62.8% | Moderate continuity |
| Skilled Birth Attendance | 312 | 62.4% | Significant leakage at delivery |
| Postnatal Care within 48 hours | 148 | 29.6% | Critical drop-off post-delivery |

While ANC initiation is relatively high at 82.4%, a significant service leakage occurs at the point of delivery, where skilled birth attendance drops to 62.4%, as tracked in Table 1. The most alarming dropout is in postnatal care, where only 29.6% of women attended within 48 hours of delivery. Qualitative data contextualize this pattern: women who delivered at home or with Traditional Birth Attendants (TBAs) perceived postnatal care as unnecessary once the immediate danger of delivery had passed, a belief actively reinforced by community norms and, in some cases, by husbands.

## Predictors of Skilled Birth Attendance: The Dominant Role of Social Endorsement

Binary logistic regression identified the primary source of health information as the most statistically powerful predictor of first-trimester ANC booking, surpassing proximity to facility, economic status, and educational level. Detailed odds ratios and statistical confidence limits for each category of primary health communicator are provided in Table 2.

**Table 2.** Logistic Regression-Predictors of First-Trimester ANC Booking by Information Source (N = 500)

| Primary Information Source | OR | 95% CI | p-value |
|---|---|---|---|
| PHC Clinical Staff (Reference) | 1.00 | 1.00[1] | 0.00[1] |
| Traditional Rulers | 3.42 | 2.15 to 5.43 | < 0.001 |
| Religious Leaders | 2.89 | 1.78 to 4.69 | < 0.001 |
| Narrative Media (Edutainment) | 2.14 | 1.34 to 3.42 | 0.002 |
| Community Health Workers | 1.67 | 1.08 to 2.57 | 0.021 |

**p<0.05; p<0.01; ***p<0.001*

As explicitly modeled in Table 2, women receiving health advocacy from Traditional Rulers were 3.42 times more likely to book early for ANC than those relying solely on clinical sources. Faith leader endorsement produced a nearly threefold increase in early booking likelihood. These findings confirm the Trust Infrastructure Theory: the PHC's clinical credibility is substantially amplified when its message is transmitted through culturally authoritative channels. Narrative media (edutainment) also emerged as a significant predictor, demonstrating that even indirect, media-based stakeholder engagement produces measurable improvements in care-seeking behaviour [6, 14].

## Predictors of Skilled Birth Attendance

A separate logistic regression examined predictors of skilled birth attendance at delivery, yielding the statistical indicators displayed in Table 3.

https://doi.org/10.21522/TIJAR.2014.13.03.Art019

**Table 3.** Binary Logistic Regression: Predictors of Skilled Birth Attendance (N = 500)

| Variable | OR | 95% CI | p-value |
|---|---|---|---|
| Husband's Approval | 4.42 | 3.21 to 6.08 | < 0.001 |
| Religious Leader Support | 3.15 | 2.45 to 4.02 | < 0.001 |
| Traditional Ruler Endorsement | 3.42 | 2.15 to 5.43 | < 0.001 |
| Edutainment Exposure | 2.14 | 1.34 to 3.42 | 0.002 |
| Proximity to PHC (< 5km) | 1.85 | 1.12 to 2.95 | 0.045 |

As outlined in Table 3, husband's approval was the strongest individual predictor of skilled birth attendance (OR = 4.42; $p < 0.001$), confirming the Garrison Effect: the male partner's attitude toward hospital delivery is more determinative of the delivery outcome than physical proximity to the facility [12, 17, 24]. Traditional ruler endorsement and religious leader support also remained strongly significant, with edutainment exposure producing a meaningful though comparatively modest effect.

## Barriers to Facility-Based Delivery: The Dual-Barricade Model

Respondents were asked to identify factors that had prevented or delayed their utilization of a PHC for delivery. Results were subsequently subjected to factor analysis, which confirmed the two-cluster structure of the Dual-Barricade Model. The primary structural and socio-psychological factors identified by respondents are detailed in Table 4.

**Table 4.** Key Barriers to Facility-Based Delivery as Identified by Respondents (N = 500)

| Barrier | % Endorsing | Description |
|---|---|---|
| Financial Constraints | 68.4% | Inability to afford transport, delivery fees, and medication costs |
| Lack of Transport | 52.2% | Especially acute in riverine and rural settlements |
| Husband's Disapproval | 46.8% | Strongest socio-psychological predictor of home delivery |
| Fear of Medical Procedures | 34.2% | Particularly caesarean sections and injections |
| Negative Health Worker Attitudes | 18.6% | Perceived disrespect and dismissiveness during prior visits |

Financial constraints were the most commonly cited barrier at 68.4%, followed by transport difficulties at 52.2%, as shown in Table 4. However, husband's disapproval, which serves as a Socio-Psychological Barricade, ranked third overall and emerged as the strongest individual predictor of home delivery in logistic regression, demonstrating that the inner barricade exerts a more decisive influence on outcome than the structural one. Fear of medical procedures at 34.2% and negative health worker attitudes at 18.6% further confirm that the relationship between the community and the health system carries unresolved tensions that structural investments alone cannot address [2, 22].

## Impact of Multi-stakeholder Interventions on Health Outcomes

Communities and households exposed to coordinated MSHPI (combining edutainment, traditional ruler decrees, religious leader messaging, and PHC outreach) demonstrated significant improvements across all measured

https://doi.org/10.21522/TIJAR.2014.13.03.Art019

maternal health indicators. The proportions of distinct behavioural and awareness advancements captured post-implementation are shown in Table 5.

**Table 5.** Observed Improvements Following Multi-stakeholder Intervention (N = 500)

| Outcome Indicator | n | % | Note |
|---|---|---|---|
| Increased Facility-Based Deliveries | 342 | 68.4% | Most impactful outcome |
| Improved Community ANC Awareness | 311 | 62.2% | Driven by edutainment |
| Increased ANC Attendance (4+ visits) | 288 | 57.6% | Gatekeeper endorsement key |
| Improved Male Involvement | 194 | 38.8% | Post-edutainment series |
| Perceived Reduction in Mortality | 122 | 24.4% | Community self-report |

The most pronounced outcome collected in Table 5 was the increase in facility-based deliveries (68.4% of respondents in intervention-exposed communities), driven primarily by male behaviour change following edutainment exposure. Chi-square analysis confirmed that the association between the 'Man of the House' edutainment series and male supportive behaviour, including transport provision and financial planning for hospital delivery, was statistically significant ($p = 0.004$). The 57.6% increase in ANC attendance was most strongly linked to traditional ruler endorsement, while improved community awareness at 62.2% reflected the synergistic effect of layered communication across stakeholder channels [4, 16].

## Qualitative Thematic Findings

**Theme 1:** The Sovereignty of the Signal Key Informant Interviews with traditional rulers, religious leaders, and PHC administrators consistently surfaced the theme of messenger credibility as the primary determinant of message adoption. Respondents described a hierarchy of trusted voices in which the clinical health worker ranked lower than the King, the Imam, or the Pastor. One PHC facility head in Ibeju Lekki noted that attendance at ANC registration events rose sharply following a formal decree by the community's traditional ruler, without any change in the services offered or the facility's physical accessibility. This pattern was termed the 'Sovereignty of the Signal', representing the point at which a health message is accepted as authoritative because of who delivered it, not what it contained.

**Theme 2:** The Garrison Effect Focus Group Discussions with male community members revealed that men did not perceive themselves as barriers to maternal care; rather, they understood themselves as protectors exercising informed caution. Several participants expressed concern about the perceived impersonal or disrespectful treatment their wives might receive at public health facilities, which had informed their preference for TBA-assisted home births. Critically, the 'Man of the House' edutainment series, which framed hospital delivery as an act of responsible fatherhood, was cited by multiple participants as having changed their perspective. This finding validates the proposition that male behaviour change requires narrative reframing, not clinical argumentation.[4,12]

**Theme 3:** Structural Fragmentation as a Systems Failure Traditional rulers and religious leaders across both LGAs independently described the same structural problem: they are rarely consulted by health authorities before health campaigns are launched. PHC managers similarly reported receiving no institutional framework for community engagement. The result is that health messaging from official channels competes with, rather than aligns with, messages from community authority structures, creating confusion and reducing uptake. This theme directly validates the Dual-Barricade Model's emphasis on systemic

fragmentation as a contributor to socio-psychological barriers.

## Discussion

The findings of this study make a compelling empirical case for a fundamental re-conceptualization of maternal health promotion in peri-urban Nigeria. The dominant paradigm, which measures success by the construction of facilities, the training of clinical staff, and the distribution of health education materials, is necessary but demonstrably insufficient in communities where social legitimacy is the primary currency of health-seeking behaviour [7].

The Trust Infrastructure Theory developed in this study provides a conceptual bridge between public health science and community sociology. The data demonstrates that clinical credibility is not intrinsic to the PHC; it is relational. It must be cultivated through deliberate engagement with the authority systems that communities have already invested with their trust. When traditional rulers endorse the PHC, they are not simply repeating a health message; they are lending their social capital to the health system, enabling it to operate within the community's existing trust economy [5].

The Dual-Barricade Model similarly advances the field by demonstrating that infrastructural improvements, while important, have limited effect on maternal mortality when the inner, socio-psychological barricade remains intact. A paved road to a PHC is irrelevant if a woman lacks her husband's permission to travel it. This finding has profound implications for resource allocation in Nigerian public health: it suggests that investments in community engagement and narrative communication should be treated as essential infrastructure, not as supplementary activities [6, 14].

The magnitude of the Garrison Effect, demonstrated by the OR of 4.42 for husband's approval as a predictor of skilled birth attendance, underscores a gap that the field has long identified but rarely addressed with structural solutions. Edutainment represents a cost-effective, culturally congruent approach to closing this gap. The 68.4% improvement in facility-based deliveries among edutainment-exposed communities is not a marginal gain; it is a transformative one, achieved not through clinical investment but through storytelling [4, 17, 24].

The postnatal care dropout rate at 29.6% remains the most critical unresolved challenge identified in this study. While the MSHPI improved ANC attendance and facility delivery rates substantially, it did not produce a proportionate improvement in postnatal engagement. This suggests that the Trust Infrastructure required for postnatal care engagement requires a distinct, targeted intervention, one that addresses the community belief that medical care ends at delivery [1, 20].

## Conclusion

This doctoral study establishes that maternal and infant survival in Lagos State is a function not of clinical investment alone, but of Integrated Advocacy: a coordinated alignment of medical expertise, traditional authority, religious legitimacy, and male household engagement that creates the social conditions for health-seeking behaviour to flourish. The Trust Infrastructure Theory and Dual-Barricade Model introduced in this study contribute original frameworks to the public health literature, providing both diagnostic tools for understanding community health barriers and prescriptive tools for designing interventions that can overcome them. The Community Health Advocacy Council (CHAC) model proposed in this study offers a scalable, institutionalizable mechanism for embedding multi-stakeholder collaboration into the routine architecture of PHC governance in Nigeria. Maternal mortality in Nigeria is not an unsolvable problem. It is a coordination problem, one that can be addressed when the

clinic, the throne, the pulpit, and the household operate not as separate systems delivering competing messages, but as a unified advocacy architecture, each amplifying the other's signal in the interest of every expectant mother's survival.

### Recommendations

#### For the Lagos State Government and Ministry of Health

Institutionalize Community Health Advocacy Councils (CHACs) in every rural and peri-urban LGA, mandating quarterly coordination between PHC administrators, traditional rulers, and clergy as a formal component of PHC governance.

Adopt and scale the narrative media model, specifically the 'Man of the House' approach, as the official framework for male engagement in maternal and reproductive health campaigns across Lagos State.

Commission a state-wide Trust Infrastructure Assessment to map the social authority structures in each LGA and identify the most credible channels for maternal health communication.

#### For Traditional and Religious Institutions

Integrate maternal health reporting as a standing agenda item in traditional ward meetings, making ANC registration and facility delivery a matter of community responsibility endorsed by royal authority.

Designate monthly 'Safe Motherhood' sessions in religious gatherings, utilizing the authority of the pulpit to dismantle misinformation regarding immunizations, caesarean sections, and hospital births.

#### For Further Research

Longitudinal studies should evaluate the sustained impact of CHACs on maternal mortality rates over a 3-to-5-year period.

Future research should investigate the role of digital platforms, including WhatsApp-based edutainment and community radio, in extending the reach of multi-stakeholder messaging in areas where in-person engagement is constrained by geography.

A comparative study across southern Nigerian states would assess the transferability of the Trust Infrastructure framework beyond Lagos.

## Ethical Approval

Ethical approval for this study was formally obtained from the Lagos State Health Research Ethics Committee under Reference Number LSHSC/HREC/24/04/53. Written and verbal informed consent were acquired from all participating respondents prior to engagement, with absolute anonymity and data confidentiality guaranteed throughout the research cycle.

## Data Availability

The datasets generated and analyzed during the study are not publicly available due to participant privacy protections and institutional ethical constraints. De-identified datasets may be made available by the corresponding author upon qualified request for verified academic research purposes, subject to appropriate institutional ethical clearance.

## Author Contributions

The sole author (B.E.O.) designed the study architecture, developed the Multidimensional Maternal Health Questionnaire (MMHQ), coordinated the field research teams during concurrent quantitative and qualitative data collection, executed the quantitative statistical analyses (including binary logistic regression and Chi-square tests) in SPSS, performed qualitative thematic content modeling, and compiled the complete original manuscript and its subsequent revisions.

## Funding

This research received no external funding or financial grants from public, commercial, or non-profit sector entities; it was entirely self-funded by the author.

## Acknowledgements

The author is grateful to the Lagos State Health Research Ethics Committee for ethical clearance, to the traditional rulers and religious leaders of Ibeju Lekki and Epe for their participation and cooperation, and to the ten field research assistants whose professionalism ensured the integrity of data collection. Special appreciation is extended to the supervisory faculty at Texila American University.

## Conflict of Interest

The author declares no conflict of interest.

## References

[1]. Afulani PA, Buback L, Kelly AM, Kirumbi L, Cohen CR, Lyndon A. Providers' perceptions of communication and women's autonomy during childbirth: A mixed-methods study in Kenya. Health Policy Plan. 2021;36(3):296-306.

[2]. Bohren MA, Vogel JP, Tunçalp Ö, Fawole B, Titiloye M, Olutayo AO, et al. Mistreatment of women during childbirth in Abuja, Nigeria: A qualitative study on perceptions and experiences of women and healthcare providers. Reprod Health. 2019;14(1):9.

[3]. Braun V, Clarke V. Using thematic analysis in psychology. Qual Res Psychol. 2006;3(2):77-101.

[4]. Doyle K, Levtov RG, Barker G, Bastian GG, Bingenheimer JB, Kazimbaya S, et al. Gender-transformative Bandebereho couples' intervention to promote male engagement in reproductive and maternal health and violence prevention in Rwanda: Findings from a randomized controlled trial. PLoS One. 2018;13(4):e0192756.

[5]. Dynes MM, Stephenson R, Rubardt M, Bartel D. The influence of perceptions of community norms on the contraceptive behavior of men and women in Ethiopia and Kenya. Health Place. 2019;18(4):1031-41.

[6]. Gabrysch S, Campbell OMR. Still too far to walk: Literature review of the determinants of delivery service use. BMC Pregnancy Childbirth. 2019;9(1):34.

[7]. Kruk ME, Gage AD, Arsenault C, Jordan K, Leslie HH, Roder-DeWan S, et al. High-quality health systems in the Sustainable Development Goals era: Time for a revolution. Lancet Glob Health. 2018;6(11):e1196-e1252.

[8]. Lagos State Health Service Commission. Strategic guidelines for community health promotion in Lagos State. Lagos: LSHSC; 2024.

[9]. McLeroy KR, Bibeau D, Steckler A, Glanz K. An ecological perspective on health promotion programs. Health Educ Q. 1988;15(4):351-77.

[10]. National Population Commission (Nigeria), ICF. Nigeria Demographic and Health Survey 2018. Abuja: National Population Commission and ICF; 2019.

[11]. Nigeria National Population Commission. Nigeria Verbal and Social Autopsy (VASA) Study on Maternal and Child Health. Abuja: National Population Commission; 2024.

[12]. Okonofua FE, Ntoimo LFC, Ogungbangbe J, Anjorin S, Imongan W, Yaya S. Predictors of men's attitude to women's use of skilled pregnancy care in rural Nigeria. BMC Pregnancy Childbirth. 2018;18(1):64.

[13]. Oligbo ME, Anya SE, Okafor IP. Comparative analysis of maternal outcomes among booked and unbooked obstetric cases in Lagos. Saudi J Med. 2024;9(2):45-53.

[14]. Oluwole EO, Adesanya T, Ige OK. Predictors of maternal healthcare services utilization among women in Lagos State. Ann Glob Health. 2025;91(1):1-12.

[15]. Olamijulo JA, Olorunfemi G, Tijani M. Trends and causes of maternal mortality at the Lagos University Teaching Hospital. BMC Pregnancy Childbirth. 2022;22(1):215.

[16]. Sacks E, Langlois ÉV, Bhattacharyya S. Community-based interventions to improve maternal health and survival: A systematic review of evidence. Bull World Health Organ. 2017;95(8):536-44.

[17]. Singh K, Brodish P, Haney E. Male involvement in antenatal care and maternal health

outcomes in Ghana and Zambia. Afr J Reprod Health. 2021;18(4):37-52.

[18]. UNICEF. Maternal and newborn health: Nigeria country data. New York: United Nations Children's Fund; 2023.

[19]. United Nations Population Fund. State of World Population 2020. New York: UNFPA; 2020.

[20]. World Health Organization. WHO recommendations on antenatal care for a positive pregnancy experience. Geneva: World Health Organization; 2016.

[21]. World Health Organization. Social determinants of health. Geneva: World Health Organization; 2019.

[22]. World Health Organization. Respectful maternity care: WHO global standards. Geneva: World Health Organization; 2022.

[23]. World Health Organization. Trends in maternal mortality 2000 to 2023. Geneva: World Health Organization; 2023.

[24]. Yargawa J, Leonardi-Bee J. Male involvement and maternal health outcomes: Systematic review and meta-analysis. J Epidemiol Community Health. 2015;69(6):604-12.